Contributions to Entomology 74 (1) 2024, 7–11 | DOI 10.3897/contrib.entomol.74.e116603

CONTRIBUTIONS TO ENTOMOLOGY

SENCKENBERG

Research Article

A new species of the genus *Rhyacophila* Pictet, 1834 (Insecta, Trichoptera, Rhyacophilidae), from India*

Tabraq Ali[1], Zahid Hussain[1], Aquib Majeed[1], Manpreet Singh Pandher[2], Sajad H. Parey[1]

1 *Insect Systematics Research Laboratory, Department of Zoology, Baba Ghulam Shah Badshah University, Rajouri, 185234, India*

2 *High Altitude Regional Centre (HARC), Zoological Survey of India, Solan, Himachal Pradesh, 173212, India*

https://zoobank.org/0087EC53-751A-4851-8E4F-AD0CA703BB87

Corresponding author: Sajad H. Parey (sajadzoo@gmail.com)

Academic editor: Simon Vitecek | Received 30 November 2023 | Accepted 19 December 2023 | Published 10 January 2024

Abstract

Rhyacophila masudi **sp. nov.** from Northwest Himalaya of India is described and illustrated. The species mostly occurs in the Pir Panjal Region of Jammu and Kashmir, India. With this new addition, the genus *Rhyacophila* is represented by 166 valid species from India.

Key Words

Asia, caddisflies, India, lactic acid, macerated

Introduction

The family Rhyacophilidae is one of the Basal Lineages of suborder Integripalpia (Morse et al. 2019) and holds significant biogeographical interest due to its primitive nature. Ross (1956) and Schmid (1970) updated the taxonomy for this family in India. These caddisflies occur primarily in north-temperate regions like North America, Europe, and northern Asia, but they can also be found in India and the tropical zones of southeastern Asia (Holzenthal et al. 2007).

In India, there are two rhyacophilid genera: *Rhyacophila* Pictet, 1834, and *Himalopsyche* Banks, 1940. *Rhyacophila* is the largest genus in the family and has more than 800 described species (Schmid 1970; Oláh 2010; Kiss 2011, 2013; Holzenthal et al. 2015; Oláh and Beshkov 2016; Valladolid et al. 2020, 2021, 2023; Morse 2023). To date, the genus *Rhyacophila* comprises 165 species and seven subspecies in India (Ali et al. 2020; Parey et al. 2023). While some species of this genus have a wide distribution in various freshwater habitats, others have a more limited range. Larvae of *Rhyacophila* species do not build cases and can move freely in the stream riverbeds. Most species have predatory larvae (Holzenthal et al. 2007), although a few feed on tiny organic particles and living vascular and nonvascular plant material (Merritt et al. 2019). The species of *Rhyacophila* occupy diverse habitats, ranging from near sea level to over 3500 m above mean sea level (AMSL), and can be found in water with different flow velocities. The majority of the Indian species of *Rhyacophila* are endemic to the Himalayan region. They prefer clean, fast-flowing fresh waters that are not affected by organic pollution. The Himalayan region in India, Pakistan, and Nepal provides a suitable environment for this genus (Parey et al. 2023).

* The paper is part of 17th International Symposium on Trichoptera, Lunz am See (Austria), 6–10 September 2021

Copyright *Tabraq Ali et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Methods

Sampling area

The adults of this species were collected from Jammu and Kashmir and adjoining localities in Himachal Pradesh, northwestern Himalaya. It has been noted from the study that the species is predominant in the Pir Panjal region of Jammu and Kashmir Himalaya. A total of 14 male specimens were collected from sites at 1770–3680 m AMSL (Figs 1, 2, Table 1).

Morphological study

Samples were collected from streams and waterfalls at high altitudes using a 15-watt ultraviolet light (BioQuip Products Inc.), powered by a sealed rechargeable 12-volt battery, and hand-picked from a white sheet and nearby stones for 1–3 hours after dusk. The collected specimens were preserved in 96% ethanol. The male genitalia were carefully extracted using fine-tipped forceps and macerated using the lactic acid technique described by Blahnik et al. (2007) and, for better visualization, we placed them in 70% ethanol with a drop of glycerine. The nomenclature for the genitalia follows that of Schmid (1970). The type specimens are stored in the Museum of the Department of Zoology, Baba Ghulam Shah Badshah University, Rajouri, Jammu & Kashmir, India (BGSBU). Variations of the morphological characteristics of the genitalia of the newly discovered species, *Rhyacophila masudi* sp. nov., were examined by using 14 male specimens. Illustrations were created using Adobe Illustrator (ADOBE CREATIVE CLOUD, version 2021) based on photographs captured with a Realme 6 Pro camera mounted on a stereozoom microscope (Olympus SZX10). The photographs and plates were assembled using Adobe Photoshop.

Results

Species description

***Rhyacophila masudi* sp. nov.**

https://zoobank.org/9E4E0165-737F-44F4-89CB-6054EFEB793E

Diagnosis. The new species resembles *Rhyacophila obscura* Martynov, 1927, in the *R. obscura* Species Group. The characters that place the new species in the

Table 1. Sampling sites, their coordinates, and elevations (m a.s.l.) for *Rhyacophila masudi* sp. nov.

Sampling Site	Region	GPS coordinates (Latitude, Longitude)	Altitude	Date of Collection
Buffliaz	Jammu & Kashmir	33.5937, 74.3647	1850 m	15.vi.2019
Kandra	Jammu & Kashmir	33.3845, 74.5976	1820 m	17.ix.2020
Kousernag	Jammu & Kashmir	33.5008, 74.8347	3680 m	12.viii.2020
Thajawas	Jammu & Kashmir	34.2921, 75.2686	2770 m	2.vii.2021
Panchpulla	Hiamchal Pradesh	32.5261, 75.9894	2000 m	28.v.2022
Bhagsunag	Hiamchal Pradesh	32.2461, 76.3369	1770 m	30.v.2022



Figure 1. Map showing the collection sites of *Rhyacophila masudi* sp. nov.



Figure 2. Pictures showing some localities of *Rhyacophila masudi* sp. nov. across different locations in the Indian Himalaya. (Abbreviations**:** J&K = Jammu and Kashmir, HP = Himachal Pradesh).

R. obscura Group are features of segment X, the aedeagus, parameres, ventral lobes of the aedeagus, and the second segment of each inferior appendage. In lateral view, segment X is short and bent downwards in *R. masudi* sp. nov. but in *R. obscura*, it is long and directed posterad. The phallotheca is crescentic in the former and rectangular in the latter species. The excision between the two lobes of the second segment of the inferior appendages is wider in *R. masudi* sp. nov. but it is shallow in *R. obscura*. In *R. masudi* sp. nov. parameres are apically rounded but oval in the latter (Fig. 4A).

Description. Adult (male): Head, thorax, and legs yellowish brown; general body pale brown and antennae with alternating light and dark brown bands. Abdomen dark brown and genitalia pale brown. Length of each antenna 4.3 mm and of each forewing 6.79 mm (n = 14). In forewings, fork II arises slightly more nearly basal than fork I (Fig. 3A–C).

Genitalia (♂) (Fig. 4). In lateral view, tergum of segment IX slightly longer than its sternum. In dorsal view, segment X trilobed, outer lobes much broader than median lobe and apicolaterally round, tiny middle lobe triangular; in lateral view bent downwards, with blunt apex. Pair of dark anal



Figure 3. Adult male of *Rhyacophila masudi* sp. nov. and picture of M. Choudhary. **A.** *R. masudi*, habitus, left lateral; **B.** *R. masudi*, right forewing, dorsal; **C.** right hind wing, dorsal; **D.** picture of the late Masud Choudhary.



Figure 4. *Rhyacophila masudi* sp. nov. male genitalia. **A.** Left lateral; **B.** Segment X, dorsal; **C.** Segment X and anal sclerites, ventral. Abbreviations: AB = apical band; AED = aedeagus; AS = anal sclerite (paired); END = endotheca; INF = inferior appendage (paired); IX = abdominal segment IX; PAR = paramere (paired); PHAL = phallotheca; TB = tergal band; VLA = ventral lobe of aedeagus; X = abdominal segment X.

sclerites elliptical in ventral view. Phallotheca crescentic in lateral view, with convex anterior margin and concave posterior margin; endotheca shorter, tubular; aedeagus slender, tubular, acute distally; ventral lobe longer than aedeagus, apically acute, directed posterodorsad; parameres elliptical, round apically, clothed with dense setae. Inferior appendages each with first segment longer than second; second segment with dorsal lobe apically round and elevated, ventral lobe slender and projecting beyond dorsal lobe, with tall, deep excision between lobes. In ventral view first segment of inferior appendage with mesal concavity.

Female. Unknown.

Type material. ***Holotype.*** ♂, INDIA: Jammu and Kashmir, Buffliaz, DKG forest, 33.5937°N, 74.3646°E, 1850 m, 15.vi.2019, Coll. Tabraq Ali, Zahid Hussain, Aquib Majeed, and Osman Javid.

Paratypes. INDIA: (2♂) same locality and collection data as holotype except 2.ix.2019; (2♂) Jammu and Kashmir, Rajouri, Kandra, 33.3844°N, 74.5975°E, 1820 m, 17.ix.2020; (2♂) Jammu and Kashmir, Kousernag, 33.5007°N, 74.8347°E, 3680 m, 12.viii.2020; (3♂) Jammu and Kashmir, Thajawas Glacier, 34.2920°N, 75.2686°E, 2780 m, 2.vii.2021; (3♂) Himachal Pradesh, Panchpulla waterfall, 32.5260°N, 75.9893°E, 2000 m, 28.v.2022; (1♂) Himachal Pradesh, Bhagsunag waterfall, 32.2460°N, 76.3369°E, 1770 m, 30.v.2022. **Collectors:** Tabraq Ali, Zahid Hussain, Aquib Majeed, and Sajad H. Parey.

Holotype and paratypes depository. Holotype and paratypes are deposited in the museum of the Department of Zoology, BGSB University, Rajouri, India.

Distribution. India: Jammu and Kashmir region, Himachal Pradesh state.

Etymology. This species is named in honor of the late Masud Choudhary (Fig. 3D), founding Vice Chancellor of Baba Ghulam Shah Badshah University, Rajouri, J&K, India, for his great contributions towards making the university a hub of excellence in teaching and research and for his excellent services to improve social and educational opportunities for the people of the Pirpanjal Range of Jammu and Kashmir.

Acknowledgements

We thank Dr Karl Kjer for the ultraviolet light used for the collection of samples. The corresponding author is thankful to the Science and Engineering Research Board (SERB), Government of India, for providing a financial grant vide sanction order no. CRG/2020/001494 Dated 1-12-2020 and providing a travel grant under ITS (International Travel Support) scheme with file no. ITS/2022/001567 dated 27 July 2022, to attend the XVII International Symposia on Trichoptera at Lunz am See, Austria. Thanks also to Dr Simon Vitecek for providing the best possible facilities to the corresponding author at Lunz am See, Austria. Thanks to Dr John C. Morse for providing the relevant literature for the current studies. We also thank anonymous reviewers for critically reading this manuscript. We highly appreciate the technical support provided by Mr Shahid Hamid Qureshi (Department of School Education, J&K) and Dr Muzaffar A. Kichloo (Department of Higher Education, J&K) during our research program.

References

Ali T, Parey SH, Pandher MS, Saini MS (2020) Checklist of the caddisfly family Rhyacophilidae (Insecta: Trichoptera) in India. Insecta Mundi 0809: 1–17. https://doi.org/10.1201/9781003410959-4

Banks N (1940) Report on certain groups of neuropteroid insects from Szechwan, China. Proceedings of the United States National Museum 88(3079): 173–220. https://doi.org/10.5479/si.00963801.88-3079.173

Blahnik J, Holzenthal RW, Prather L (2007) The lactic acid method for clearing Trichoptera genitalia. In: Bueno-Soria J, Barba-Alvarez R, Armitage BJ (Eds) Proceedings of the 12th International Symposium on Trichoptera, 18–22 June 2006. The Caddis Press, Columbus, Ohio, 9–14.

Holzenthal RW, Blahnik RJ, Prather AL, Kjer KM (2007) Order Trichoptera Kirby, 1813 (Insecta), caddisflies. Zootaxa 1668: 639–698. https://doi.org/10.11646/zootaxa.1668.1.29

Holzenthal RW, Thomson RE, Rios-Touma B (2015) Order Trichoptera. In: Thorp JH, Rogers C (Eds) Thorp and Covich's Freshwater Invertebrates: Ecology and General Biology, 4th Edn., volume 1. Academic Press, 965–1002. https://doi.org/10.1016/B978-0-12-385026-3.00038-3

Kiss O (2011) Two new species of *Rhyacophila* (Trichoptera, Rhyacophilidae) from Nepal. Acta Zoologica Academiae Scientiarum Hungaricae 57(2): 111–116. https://doi.org/10.11646/zootaxa.2991.1.8

Kiss O (2013) Three new species of *Rhyacophila* (Trichoptera: Rhyacophilidae) from Taiwan and Nepal. Zootaxa 3640: 213–223. https://doi.org/10.11646/zootaxa.3640.2.5

Martynov AV (1927) Contributions to the aquatic entomofauna of Turkestan. I. Trichoptera Annulipalpia. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de Saint Pétersbourg 28: 162–193.

Merritt RW, Cummins KW, Berg MB (2019) An Introduction to the Aquatic Insects of North America, 5th Edn. Kendall Hunt Publishing Company, Dubuque, Iowa, 1480 pp.

Morse, JC, Frandsen PB, Graf W, Thomas JA (2019) Diversity and ecosystem services of Trichoptera. In: Morse JC, Adler PH (Eds) Diversity and Ecosystem Services of Aquatic Insects. Insects 10: 125. https://doi.org/10.3390/insects10050125

Morse JC [Ed.] (2023) Trichoptera World Checklist. https://trichopt.app.clemson.edu/ [accessed 07 December 2023]

Oláh J (2010) New species and new records of Palearctic Trichoptera in the material of the Hungarian Natural History Museum. Annales Historico-naturales Musei Nationalis Hungarici 102: 65–117.

Oláh J, Beshkov S (2016) New records of Trichoptera in the Balkan Peninsula and Romania, with description of new *Rhyacophila* sibling species by speciation traits. Folia Historico-Naturalia Musei Matraensis 77: 87–104. https://doi.org/10.17112/FoliaEntHung.2016.77.87

Parey SH, Ali T, Pandher MS (2023) Rhyacophilidae of India (1st ed.). Jenny Stanford Publishing, Singapore, 320 pp. https://doi.org/10.1201/9781003410959

Pictet FJ (1834) Recherches pour servir à l'histoire et l'anatomie des Phryganides. A. Cherbuliez, Geneva, 235 pp. [20 plates] https://doi.org/10.5962/bhl.title.8547

Ross HH (1956) Evolution of the Mountain Caddisflies. University of Illinois Press, Urbana, Illinois, 213 pp.

Schmid F (1970) Le genre *Rhyacophila* et la famille des Rhyacophilidae (Trichoptera). Mémoires de la Société Entomologique du Canada 66: 1–230. https://doi.org/10.4039/entm10266fv

Valladolid M, Kučinić M, Arauzo M, Cerjanec D, Ćuk R, Dorda BA, Lodovici O, Stanić Koštroman S, Vučković I, Rey I (2020) The *Rhyacophila fasciata* Group in Croatia and Bosnia and Herzegovina: *Rhyacophila f. fasciata* Hagen, 1859 and the description of two new subspecies, *Rhyacophila fasciata delici* Kučinić & Valladolid (ssp. nov.) from Croatia and Bosnia and Herzegovina and *Rhyacophila fasciata viteceki* Valladolid & Kučinić (ssp. nov.) from Bosnia and Herzegovina (Trichoptera: Rhyacophilidae). Zootaxa 4885(1): 051–075. https://doi.org/10.11646/zootaxa.4885.1.3

Valladolid M, Arauzo M, Chertoprud M, Chvojka P, Czachorowski S, Dorda BA, Hinić J, Ibrahimi H, Karaouzas I, Krpač V, Kučinić M, Lodovici O, Salokannel J, Slavevska-Stamenković V, Stojanović K, Wallace I, Rey I (2021) The *Rhyacophila fasciata* Group in Europe: *Rhyacophila fasciata* Hagen 1859 and formerly synonymized species (Trichoptera: Rhyacophilidae), with new description of *Rhyacophila fasciata* and *Rhyacophila septentrionis* McLachlan 1865 (stat. prom.). Zootaxa 4975(1): 1–57. https://doi.org/10.11646/zootaxa.4975.1.1

Valladolid M, Waringer J, Arauzo M, Chvojka P, Dorda BA, Komzák P, Lodovici O, Rey I (2023) The *Rhyacophila fasciata* Species Complex (Trichoptera: Rhyacophilidae) in Central Europe with description of a new species, *Rhyacophila loeffleri* Valladolid & Waringer, n. sp., based on morphological, genetic and ecological evidence. Zootaxa 5325: 451–482. https://doi.org/10.11646/zootaxa.5325.4.1